## Drug Facts

## Active Ingredients (in each tablet) Purpose Aluminum hydroxide dried gel 200 mg . Magnesium hydroxide 200 mg ..... Antacid .Antacid Simethicone 25 mg. Antigas

## Uses Relieves:

• Heartburn • Sour stomach • Acid indigestion • Bloating, pressure and discomfort commonly

## Warnings

Ask a doctor before use if you have • Kidney disease • A magnesium restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids may interact with certain

Stop use and ask a doctor if symptoms last for more than 2 weeks If pregnant or breast-feeding, ask a health professional before use.

Keep out of the reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away 1.800.222.1222

Directions • Adults and children 12 years of age and older: Chew 2 to 4 tablets 4 times daily. Repeat hourly if symptoms return, or as directed by a physician.

• Children under 12: ask a doctor

- Do not take more than 12 tablets in 24-hour period, or use the maximum dosage for more than 2 weeks, except under the advice and supervision of a physician

**Other Information** • Each tablet contains: magnesium 95 mg • Store at 59° to 77°F (15° to 25°C)

■ Tamper Evident. Do not use if blister or printed Gelusil foil seal is broken

See end panel for lot number and expiration date

Inactive ingredients Dextrose, flavors, magnesium stearate, maltodextrin, silica, sucralose

Questions? Call 1-844-241-5454

©2016 WellSpring Sarasota, FL 34243 USA